

RESEARCH

Chemosensory Pleasure Scale: Validity and Reliability of Turkish Version

Kimyasal Duyu Keyif Ölçeği: Türkçe Versiyonunun Geçerliliği ve Güvenirliği

Özge Akgül¹ , Betül Önder Uzgan² 

Öz

Bu çalışmanın amacı, tat ve koku almada hedonik hazı belirlemek için geliştirilen Kimyasal Duyu Keyif Ölçeği'nin Türkçe versiyonunun geçerlilik ve güvenilirliğini değerlendirmektir. Bu çalışma 247 katılımcı ile gerçekleştirilmiştir. Verilerin toplanmasında Sosyodemografik Form (SF), Kimyasal Duyu Keyif Ölçeği-Türkçe Versiyonu (CPS-TR), Fiziksel Anhedoni Ölçeği (FAÖ) ve Sosyal Anhedoni Ölçeği (SAÖ) kullanılmıştır. Güvenirlik analizi için Cronbach alfa iç tutarlılık katsayısı, madde-toplam puan korelasyon katsayıları ve Spearman Brown split-half testi hesaplanırken, ölçeğin yapı geçerliliğini tahmin etmek için açıklayıcı faktör analizi yapılmıştır. Ölçeğin iç tutarlılık güvenirliliği dikkate alındığında, Cronbach alfa güvenirlilik katsayısı $\alpha = 0.87$ olarak bulunmuştur. Ölçek maddelerinin madde ve toplam puan korelasyon katsayıları 0.35 ile 0.70 arasında bulunmuştur. Açıklayıcı faktör analizi toplam varyansın %68.9'unu açıklamış ve ölçeğin üç faktörlü yapısını desteklemiştir. Bu faktörlerin faktör yükleri 0.55 - 0.85 arasında tahmin edilmiştir. CPS-TR'nin FAÖ ve SAÖ ile korelasyonu istatistiksel olarak anlamlı bulunmuştur. CPS ölçeği Türkçe dilinde kimyasal hazı değerlendirmek adına geçerli ve güveniliridir.

Anahtar sözcükler: Kimyasal duyu keyif ölçeği, geçerlilik, güvenirlilik

Abstract

The aim of this study was to assess the validity and reliability of the Turkish version of the Chemosensory Pleasure Scale, which is developed to determine the hedonic pleasure in taste and smell. The present study was conducted with a sample of 247 participants. Sociodemographic Form (SF), Chemosensory Pleasure Scale Turkish Version (CPS-TR), Physical Anhedonia Scale (PAS), and Social Anhedonia Scale (SAS) were used to collect data. For the reliability analysis, Cronbach alpha internal consistency coefficient, item-total score correlation coefficients, and Spearman-Brown split-half test were calculated, whereas Explanatory factor analysis was performed to estimate the construct validity of the scale. Considering the internal consistency reliability of the scale, Cronbach's alpha reliability coefficient was found to be $\alpha=0.87$. The item and total score correlation coefficients of the scale items were found between 0.35 and 0.70. Exploratory factor analysis explained 68.9% of the total variance and supported the three-factor structure of the scale. Factor loadings of these factors were estimated between 0.55-0.85. The correlation of CPS-TR with PAS and SAS was statistically significant. The results of this study suggest that the Turkish version of the CPS is a reliable and valid instrument for determining chemosensory pleasure.

Keywords: Chemosensory pleasure scale, validity, reliability

¹ İzmir Democracy University, İzmir, Turkey

² Çanakkale Onsekiz Mart University, Çanakkale, Turkey

✉ Özge Akgül, İzmir Democracy University, Faculty of Arts and Sciences, Department of Psychology, İzmir, Turkey
ozge.akgul@idu.edu.tr | 0000-0001-8321-5485

Received: 01.04.2021 | Accepted: 15.05.2021 | Published online: 28.06.2021

ANHEDONIA is characterized as a diminished interest in likable events or activities, including sensory, social, or behavioral experiences (Nusslock and Alloy 2017). Loss of pleasure impairs quality of life and may affect social functioning (Miller 1987). Considering experiencing pleasure is an essential part of well-functioning lives, anhedonia exists in many neuropsychiatric disorders like schizophrenia, depression, addiction, and Alzheimer's disease (Gard et al. 2007, McCabe 2014). Anticipatory (wanting) and consummatory (liking) pleasure are the two components of hedonic experience; while the former reflecting the anticipation of rewarding stimulus, the latter refers to the reaction to in the moment pleasure (Kring and Elis 2013). Neuroscientific studies showed that there are two different pathways in our brains regarding these two components (Berridge and Robinson 1998, Berridge 2009). In order to evaluate these domains in pleasure experience, several self-report scales have been developed, such as the Temporal Experience of Pleasure Scale (Gard et al. 2007, Chan et al. 2010), the Anticipatory and Consummatory Interpersonal Pleasure Scale (Gooding and Pflum 2014), the Chapman Physical Anhedonia Scale (CPAS), the Chapman Social Anhedonia Scale (CSAS; (Chapman et al. 1976)) and the Snaith-Hamilton Pleasure Scale (SHAPS; (Snaith et al. 1995)). Of these, Turkish reliability and validity studies were published for the Chapman Physical Anhedonia Scale (Baskak et al. 2009), the Chapman Social Anhedonia Scale (Cihan et al. 2015) and the Snaith-Hamilton Pleasure Scale (Kesebir et al. 2015) and Turkish validity and reliability of the Temporal Experience of Pleasure Scale is currently an ongoing project by the writer of this article.

In addition, the ability to experience pleasure driven by taste or smell is called chemosensory hedonic capacities (Li et al. 2020). It is thought that olfaction compared to any other sensory stimuli, might be more sensitive in the evaluation of anhedonia (Li et al. 2020). Therefore, another important symptom of neuropsychiatric disorders can be thought of as olfactory anhedonia. It is known that there is a significant overlap in the brain regions like the orbitofrontal cortex (OFC), amygdala, and hippocampus concerning emotional processing and olfaction (Kamath et al. 2013). Many of the researchers have reported that olfactory anhedonia is present in depression (McCabe 2014), Parkinson's disease (Mrochen et al. 2016), anorexia nervosa (Jiang et al. 2010), and schizophrenia, in a way that both hedonic odor processing and odor identification are shown to be impaired (Moberg et al. 2014). In addition, in patients with schizophrenia, it was reported that increased negative symptoms are associated with increased olfactory anhedonia (Kamath et al. 2013).

Just like olfactory anhedonia, taste anhedonia can also be thought of as being one of the crucial features of neuropsychiatric disorders. In depression, researches showed that patients might have higher sweet taste perception (Berlin et al. 1998), lessened appetites, less desire to eat, and decreased pleasure toward food (Kazes et al. 1993). Moreover, in alcohol dependence and anorexia nervosa, abnormal hedonic taste capacities are also reported (Krahn et al. 2006). These above-mentioned scales measure several aspects of anhedonia; however, none of them evaluates chemosensory pleasure. Because these capacities may be important in understanding the nature of neuropsychiatric disorders, an assessment tool for the chemosensory pleasure scale is essential. Moreover, previous findings also proposed that patients with schizophrenia showed impairments in anticipatory pleasure but not in consummatory pleasure (Gard et al. 2007, Yan et al. 2019). However, it is unknown whether patients show the same pattern in chemosensory hedonic experience or not. And this is another critical reason to develop a valid self-

report scale which helps to reveal both anticipatory and consummatory pleasure aspects in order to understand taste and smell hedonic responses.

All these findings suggested that olfactory anhedonia may be a significant indicator of neuropsychiatric disorders, and there were no practical tools for evaluating chemosensory pleasure. Therefore in 2019, the Chemosensory Pleasure Scale (CPS) was developed in order to evaluate taste and smell hedonic pleasure in order to understand various disorders (Zhao et al. 2019). Considering CPS is a new, practical assessment tool, we have aimed to study the psychometric properties of the Chemosensory Pleasure Scale to be used in the Turkish population. CPS is an instrument that was developed to determine the hedonic pleasure in taste and smell. The aim of the study was to translate CPS into Turkish and study the reliability and validity of the scale. The permission has been obtained from the developer of the scale.

Method

Sample

The inclusion criteria in the study included being between the ages of 18-55, being at least a primary school graduate, and having the mental capacity to fulfill the research instructions. Exclusion criteria from the study were any medical conditions that affect cognitive functions. For this reason, the participants were asked whether they had any diagnosed psychiatric disorders or not. If they had anxiety disorders, major depressive disorder, schizophrenia, bipolar disorder diagnosis, we excluded these participants. The data was obtained through an online survey between July 2020 to February 2021. 247 participants were included into study however 3 participants were excluded because they indicated that they had generalized anxiety disorders, unipolar depression and anorexia nervosa diagnosis

Measures

Sociodemographic Form (SF)

The form was developed by the researchers in order to evaluate sociodemographic characteristics of participants such as age, gender, marital status, and educational status, as well as the questions about the patterns of alcohol, nicotine, and substance use.

Chemosensory Pleasure Scale (CPS)

The scale was developed by Zhao et al., and the aim is to assess the hedonic capacity for smell and taste pleasure (Zhao et al. 2019). The Chemosensory Pleasure Scale (CPS) is a self-report scale that evaluates the enjoyment that an individual receives from stimulation of smell and taste. It includes 3 factors and consists of 12 items: with the dimensions of "consummatory", "anticipatory," and "purely olfactory," reflecting the hedonic eating, anticipating food, and smelling natural scents, respectively. Participants are expected to evaluate the pleasantness of their hedonic experience of smell and taste on a 6-point Likert scale (1 "very false for me" and 6 "very true for me"). A lower score in Chemosensory Pleasure Scale Turkish version (CPS-TR) indicates that the chemosensory anhedonia is severe. The Cronbach's alpha coefficient was 0.93 and test-retest reliability of CPS was 0.73. First factor "food" included 5 items and accounted for 46.979% of the variance. 4 items were loaded onto the second factor "imagination" with

a 14.968% of the variance and factor 3 “nature” had 3 items which accounted for 10.634% of the variance.

Physical Anhedonia Scale (PAS)

Physical Anhedonia Scale was developed by Chapman et al. in 1976 (Chapman et al. 1976). Physical anhedonia includes physical experiences such as food, touch, sexuality, warmth, movement, smell, and sound. Baskak et al. reported in 2009 that 50 out of 61 items are valid and reliable in the Turkish population (Baskak et al. 2009).

Social Anhedonia Scale (SAS)

The scale was first created by Chapman et al. in the form of a 48-item self-report scale to measure social satisfaction and social anxiety (Chapman et al. 1976). Mislove and Chapman reported the psychometric properties of the 40-item final version. Its Turkish validity and reliability study was conducted in 2015 by Cihan et al (Cihan et al. 2015).

Translation process

Before starting the study, the developer of the scale was contacted by the team via email, and permission for translation was obtained. The translation process was independently carried out by two independent psychologists with an advanced degree in English. After this procedure, researchers agreed on the best version of the translation; items thought to best express each item have been determined and it was compared to the original form by the same team. The back-translation was applied by another researcher who is also advanced in English. and The back-translated questionnaire was sent to the developer of the instrument, and feedback was obtained for the accuracy and clarity of the translation.

After these phases, a 12-item Turkish form was obtained. Izmir Democracy University Social Sciences and Humanities Research and Publication Ethics Committee (7.7.2020 date with the number of 2020/09-06) has approved the study prior to the start of the study. For evaluating the applicability and comprehensibility of items of CPS-TR, 58 participants (above 18 years, 53 females, 5 males) were administered a pilot study online. After feedback of participants, most understandable and appropriate sentences for each item were used in the Turkish form of original scale in this study. The last version of CPS-TR was applied to participants. The rules of the Declaration of Helsinki were complied with during the data collection process. The final form of the questionnaire was applied to the participants with a consent form, Sociodemographic Form (SF), Chemosensory Pleasure Scale Turkish version (CPS-TR), Physical Anhedonia Scale (PAS), and Social Anhedonia Scale (SAS). In the exploratory factor analysis, it was known that at least 5-10 participants per item were required. Therefore a total of 247 participants were enrolled in the study (Sencan 2005).

Statistical analysis

Cronbach alpha internal consistency coefficient, item-total score correlation coefficients, and Spearman-Brown split-half test were calculated for reliability analysis. Exploratory factor analysis was performed to estimate the construct validity of the scale. Items with a factor load over 0.4 were included in the evaluation. The convergent validity of CPS-TR was assessed by conducting a Pearson correlation. Statistical

analyses were made with SPSS Statistics 20.0. The level of significance was set at $p < 0.05$.

Results

Among the participants, 59 (23.9%) were male, and 187(75.7%) were female, and 1 was (0.4%) queer with a mean age of 32.62 ± 11.27 . The sociodemographic characteristics of the participants were summarized in Table 1.

Table 1. Sociodemographic characteristic of participants

Variables	Mean \pm SD	
Age	32.62 \pm 11.27	
Total Year of Education	17.17 \pm 3.48	
Gender	N (%)	
Female	187(75.7%)	
Male	59 (23.9%)	
Queer	1 (0.4%)	
Education		
University	229 (92.7%)	
Highschool	17 (6.9%)	
Elementary School	1 (0.4%)	
	Yes (%)	No (%)
Smoking	70.9%	29.1%
Alcohol	70.9%	29.1%
Other Substance use(drugs etc.)	4.9%	95.1%

Table 2. CPS-TR item total correlation and Cronbach's alpha

Items	Corrected Item-total correlation	Cronbach- α if item deleted
Item1	0.353	0.875
Item2	0.588	0.863
Item 3	0.423	0.872
Item4	0.615	0.861
Item5	0.428	0.873
Item6	0.569	0.866
Item7	0.667	0.858
Item8	0.700	0.855
Item9	0.633	0.860
Item10	0.641	0.861
Item11	0.668	0.857
Item12	0.547	0.868

F1: Imagination: ability to experience pleasure when guessing food (Items: 2,4, 7, 8, 11); F2:Food: ability to experience pleasure while eating food (Items: 6, 9, 10, 12); F3: Nature: ability to experience pleasure when smelling natural scents (Items: 1, 3, 5)..

Reliability analysis

Cronbach alpha was used to calculate internal consistency coefficient of the scale was found as 0.87. Literature indicates Cronbach alpha levels as follows; excellent (0.93–0.94), strong (0.91–0.93), reliable (0.84–0.90), robust (0.81), fairly high (0.76–0.95), high (0.73–0.95), good (0.71–0.91), relatively high (0.70–0.77). Our result showed that cronbach alpha 0.87 reflected high reliability of the scale (Taber 2018). Corrected item-total score correlations and Cronbach's alpha values when the item is deleted are shown

in Table 2. In this study, we did not delete any of the items in the original scale because Cronbach alpha value did not increase. Item total correlation of all items in the scale were bigger than .30. Then, the split-half method was performed. The Spearman-Brown coefficient was found as 0.78 which means reliability of scale is at a good level (Taber 2018).

Table 3. Results of exploratory factor analysis (EFA)

Items	F1	F2	F3	Communalities
1) Yağmur sonrası, çimenle karışmış toprak kokusu beni mutlu eder.			0.817	0.694
2)Televizyonda pişirilen iyi bir yemek gördüğümde neredeyse kokusunu alabilirim	0.760			0.619
3)Yürüyüş için dışarı çıktığımda temiz havayı derince solumanın verdiği histen hoşlanırım.			0.784	0.656
4)Her yemek kokusu aldığımda mutlu hissederim	0.555			0.492
5)Yağmur sonrası parkta temiz havayı soluyabilmeyi dört gözle beklerim.			0.809	0.726
6)Lezzetli yiyecekler beni mutlu eder.		0.854		0.769
7)Bir yiyecek reklamı gördüğümde, o yiyeceğin tadını hayal etmeye başlarım	0.805			0.711
8)Bir restoranda menüyü incelerken yiyeceğin tadını hayal etmeyi severim.	0.826			0.755
9)Lezzetli yiyecekler yemek dört gözle beklenen bir şeydir.		0.791		0.750
10)Yemek yerken iyi hissederim.		0.782		0.729
11)En sevdiğim yiyeceği hayal ettiğimde lezzetini neredeyse tadarım.	0.744			0.652
12)En sevdiğim yemeği yemekten zevk alırım.		0.826		0.718
Variance Explained (%)	26.30	25.11	17.48	68.918
Eigenvalue	5.225	1.749	1.295	
Cronbach alpha	.85	.86	.75	

Validity analysis

To assess whether the scale is appropriate for factor analysis, first we performed Keiser Meyer Olkin and Barlett sphericity test. The Kaiser Meyer Olkin coefficient was calculated, and the sample was found to be suitable for factor analysis. The KMO sample fit coefficient was 0.87, which falls into the range of being good. The correlation between variables was sufficient according to the Barlett sphericity test ($\chi^2 = 1452.903$, $df = 66$, $p < 0.001$). Therefore, it was appropriate for factor analysis. After the Varimax rotation technique was applied, items with an eigenvalue greater than 1 and factor loadings greater than 0.40 were evaluated. It is recommended for 200 participants, factor loads should be greater than 0.364 (Field A 2009). In the light of these results, none of the items of scale were excluded. Then, Three factors were obtained as the original scale. These factors were named as 'Imagination', 'Food' and 'Nature'. These 3 factors explained 68.918% of the total variance. Factor 1, Factor 2, and Factor 3 account for 26.300%, 25.106%, 17.482% of variances, respectively. Factor loadings are explained in Table 2. Scree plot is another reliable method which is used to determine the number of factors. Point of inflexion (where the slope is flattened) occurred at the fourth factor therefore, three factors were extracted (see Figure 1), (Field 2009). Details of factor analysis were shown in Table 3.

Copyright of Current Approaches in Psychiatry / Psikiyatride Guncel Yaklasimlar is the property of Psikiyatride Guncel Yaklasimlar and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.