



ORIGINAL ARTICLE

# Translation and Psychometric Evaluation of the Turkish Version of the Motivation to Change Lifestyle for Dementia Risk Reduction Scale (T-MOCHAD-10)

Merve Aliye Akyol<sup>1,2,\*</sup>, Özlem Küçükçüçlü<sup>1,2</sup>, Burcu Akpınar Söylemez<sup>1</sup>, Bilgehan ÖZKAYA SAGLAM<sup>1,3</sup>, Déborah Oliveira<sup>4,5</sup>

<sup>1</sup> Department of Internal Medicine Nursing, Dokuz Eylul University Faculty of Nursing, Inciralti, Izmir, Turkey

<sup>2</sup> Home Care Application and Research Center, Dokuz Eylul University, Izmir, Turkey

<sup>3</sup> Internal Medicine Nursing Master Programme, Dokuz Eylul University Institute of Health Sciences, Izmir, Turkey

<sup>4</sup> Universidade Federal de São Paulo, Medical School, Division of Psychiatry, São Paulo, Brazil

<sup>5</sup> Inter American Development Bank, Division of Social Protection and Health, Washington D.C., United States

Received 21 October 2021; accepted 14 March 2022

Available online 9 June 2022

## KEYWORDS

Dementia;  
Lifestyle;  
Risk Reduction  
Behavior;  
Motivation;  
Psychometrics

## Abstract

**Background and Objectives:** The number of people living with dementia is increasing globally, particularly in middle-income countries like Turkey. To reduce risk of dementia, the implementation of lifestyle changes targeting modifiable risk factors are important. This study aimed to translate and validate the Turkish version of the Motivation to Change Lifestyle for Dementia Risk Reduction (T-MOCHAD-10).

**Methods:** Cross-sectional psychometric study design was used. After translation and back translation, we assessed face and content validity. For construct validation and reliability assessment, we conducted a survey with 601 individuals aged 40 and above using a socio-demographic form and the T-MOCHAD-10. We used Exploratory Principal Axis Factoring with Oblimin rotation to explore the factor structure. We then confirmed the factor structure using fit indices. Reliability was established using test-re-test, Cronbach's alpha coefficient, item-total and item-subdimensions correlations, ceiling and floor effects, and the Hotelling's T-squared test.

**Results:** The T-MOCHAD-10 showed adequate face and content validity (Kendall W=0.09, p=0.60). As in the original scale, a two-factor solution was obtained. All fit indices were  $\geq 0.95$ , and RMSEA was 0.06. A Cronbach's alpha of 0.79 was obtained and no significant differences were found between test and re-test measures (p>0.05). We found no evidence of ceiling or floor effects, nor response bias (Hotelling's T-squared=4683.80, p<0.001).

\* Correspondence Author: Merve Aliye Akyol, Research Assistant, PhD, Department of Internal Medicine Nursing, Dokuz Eylul University Faculty of Nursing, Inciralti, Izmir, Turkey, Home Care Application and Research Center, Dokuz Eylul University, Izmir, Turkey, Phone: 00902324124774, Mobile phone: 00905334057519

E-mail addresses: [merve.akyol@deu.edu.tr](mailto:merve.akyol@deu.edu.tr) (M.A. Akyol), [ozlem.kguclu@deu.edu.tr](mailto:ozlem.kguclu@deu.edu.tr) (Ö. Küçükçüçlü), [burcu.akpinar@deu.edu.tr](mailto:burcu.akpinar@deu.edu.tr) (B. Akpınar Söylemez), [bilgehan.ozkaya@deu.edu.tr](mailto:bilgehan.ozkaya@deu.edu.tr) (B. Özkayasaglam), [oliveiradc.phd@gmail.com](mailto:oliveiradc.phd@gmail.com) (D. Oliveira).

<https://doi.org/10.1016/j.ejpsy.2022.03.004>

0213-6163/© 2022 Asociación Universitaria de Zaragoza para el Progreso de la Psiquiatría y la Salud Mental. Published by Elsevier España, S.L. U. All rights reserved.

**Conclusion:** T-MOCHAD-10 is a short measurement tool which provides valid and reliable scores concerning the motivation to change lifestyle for reducing dementia risk among Turkish individuals aged 40 and over.

© 2022 Asociación Universitaria de Zaragoza para el Progreso de la Psiquiatría y la Salud Mental. Published by Elsevier España, S.L.U. All rights reserved.

## Introduction

The number of older people is increasing worldwide and the prevalence of diseases which are more likely to affect this population group is also rising. Dementia is a neuropsychiatric disorder caused by a variety of brain illnesses such as Alzheimer's disease and vascular dementia and is one of the most disabling and burdensome diseases affecting older people globally.<sup>1</sup> The number of people living with dementia around the world, especially in middle-income countries like Turkey, has increased in the past decades and is expected to reach 75 million by 2030 and 132 million by 2050.<sup>2</sup> In Turkey, there were approximately 800 thousand people living with dementia in 2019, and this number is expected to increase to 3 million by 2050. Considering the anticipated growth of approximately 277% between 2019-2050, public health approaches to reduce dementia risk is crucial.<sup>3</sup> Although some known risk factors for dementia such as advanced age and genetic predisposition cannot be changed, there are many risk factors which can be modified through engaging with lifestyle changes or implementing adequate treatment to reduce risk of dementia and its global prevalence. Such risk factors include physical inactivity, unhealthy diet, harmful use of alcohol and tobacco smoking, management of weight and dyslipidemia, diabetes, and hypertension (especially in midlife)<sup>4</sup>. Building community awareness about the risk and protective factors for dementia are important, however changing lifestyle strongly depends on people's attitudes and beliefs towards dementia prevention, which will then influence the extent to which public health actions may be successful in reducing dementia incidence.<sup>5</sup> It is therefore important to understand people's levels of motivation to change lifestyle for dementia risk reduction in order to effectively reduce the number of people living with dementia globally.

The construct of motivation to change lifestyle for dementia risk reduction encompasses and/or links to several psychological and educational domains, such as dementia literacy, knowledge of dementia-related risk factors, and ways to address or reduce these; perceived susceptibility/severity of dementia (feeling under the threat of developing dementia), perceived benefits (a particular behavior change is bound to result in one not developing dementia), and self-efficacy (the extent to which one believes to be able to change an unhealthy behavior).<sup>6-9</sup> Studies show that a person's level of motivation to change lifestyle for dementia risk reduction can be influenced by individual and contextual factors, such as a family history of dementia, prior experience as a caregiver of a person living with dementia<sup>10-12</sup>, willingness to know their own risk<sup>10</sup>, and worry or fear of having dementia.<sup>13,14</sup>

Motivation is a dynamic psychological trait that can be modified during a change process or intervention.<sup>15</sup> Determining individuals' motivation for behavioral and lifestyle change for dementia risk reduction is therefore the first step to building public health strategies that can effectively

reduce dementia risk in the population. Measurement tools to evaluate individual motivation to change lifestyle for dementia risk reduction should be dementia-specific.<sup>8</sup> The Motivation to Change Lifestyle for Dementia Risk Reduction (MOCHAD-10) is a 10-item scale developed in the United Kingdom. The MOCHAD-10 is informed by five of the seven domains of the Health Belief Model<sup>16,17</sup>: self-efficacy, perceived benefits, cues to action, perceived severity, and perceived susceptibility. It is a short and robust scale, likely to be readily used in clinical practice, and validated with a large national sample of people aged 50 and above using both exploratory and confirmatory factor analyses.<sup>8</sup> This study aimed to translate the MOCHAD-10 into Turkish and to evaluate its psychometric properties in the Turkish population.

## Methods

### Design

A cross-sectional psychometric study design was employed to evaluate the validity and reliability of the Turkish version of the MOCHAD-10 scale (T-MOCHAD-10). The results were reported in line with the quality criteria for measurement properties of health status questionnaires.<sup>18</sup> The study was conducted in three sequential steps: 1) Translation and back translation of the original scale (from British English into Turkish); 2) Face and content validity of the T-MOCHAD-10, and 3) Construct validation and reliability assessment of the T-MOCHAD-10.

#### 1) Translation and back translation of the original scale

First, the permission of the scale's first author (Dr. Oliveira) was sought to translate and validate the scale for use in Turkey. The instrument was then translated from British English into Turkish by MAA, ÖK, BAS, and BÖS. As a group, the researchers then reviewed and revised each scale item for linguistic accuracy, meaningfulness, and conceptual equivalence. The translated version was then back-translated by a professional bilingual translator unfamiliar with the original version of the scale.<sup>19</sup> The back-translated form and the original form were compared by the researchers, and the lead author of the original version was also consulted and approved the final version.

#### 2) Face and content validity of the T-MOCHAD-10

Face validity check helped determine whether the items of the instrument were relevant and meaningful for the target individuals.<sup>20</sup> A group of 15 individuals aged 40 and over who did not have dementia (self-report) were invited to take part via social media. An anonymous online form was

designed by the researchers and contained the following questions: What are your general comments on the instrument? What are your comments on each of the statements? How long did you take to complete this instrument? Is the number of questions acceptable? Does the order of questions make sense to you? Did you find it difficult to answer or understand any of the items? Overall, how understandable and simple is the instrument? Have you experienced problems with the rating of the statements, and do you have any suggestions on this? Do you have any other suggestions for the instrument? Did you need help completing the instrument? If yes, who helped you and why?

Content validity was then checked by consulting nine research experts: four nursing academics (three with expertise in psychometrics and one expert in both geriatric nursing and applied psychometrics) and three physicians (one neurologist and two were geriatricians). Of the nursing academics, two experts were clinic nurses, one was a neurology nurse with more than eight years of experience working in the intensive care unit, and the other was a geriatric nurse. We asked experts to rate each item using a four-point Likert scale (1. inappropriate, 2. it should be made more appropriate, 3. it is appropriate, but needs minor changes, and 4. appropriate).

### 3) Construct validation and reliability assessment of the T-MOCHAD-10

*Setting and sample:* An anonymous online cross-sectional survey of people aged 40 and over who had no self-reported dementia diagnosis (n=601) was conducted between April-May 2021 in Turkey. Given the restrictions imposed by the COVID-19 pandemic, data were collected online. We did not specify any particular group or forum to do that and published posts regularly on WhatsApp groups, Instagram, and Facebook accounts to reach out to the community. Those who accessed the link to the online anonymous survey then received information about the eligibility criteria and accepted or not to take part. A convenience and snowballing sampling approach was used with the aim to reach a minimum of 500 participants, which would mean 50 participants per scale item to be validated (10 items in total).<sup>21</sup>

*Survey items:* The online form was designed by the researchers using the Google<sup>®</sup> platform. On the landing page, information was given about the study's aim and rationale, as well as ethical considerations of anonymity and confidentiality. The survey was then divided into two sequential parts as follows.

- 1) Socio-demographic data: age, sex, income, marital status, education level, a family member living with dementia (yes/no), caregiver of people with dementia (yes/no).
- 2) The Turkish version of the Motivation to Change Behavior for Dementia Risk Reduction Scale (T-MOCHAD-10): 10 items responded on five-point Likert-type scales (from 1. strongly disagree to 5. strongly agree). The instrument includes two subscales: positive cues to action ( items 1, 2, 3, 4, 5) and negative cues to action ( items 6, 7, 8, 9, 10). Each item is added up to calculate each subscale's score. Higher scores indicate higher motivation to change lifestyle and health behaviors to reduce dementia risk.

*Procedure:* Individuals who received the survey link through social media were provided with information about the study and were asked to confirm whether they met the eligibility criteria and agreed with the participation terms. In the end, participants were asked to write their names and e-mail addresses in case they wished to be contacted to participate in the re-test (2-3 weeks later) so we could evaluate the stability of the scale over time.<sup>22,23</sup> The survey was entirely anonymous. Participants were asked to write their name and email address only at the end of the survey and only in case they had an interest in taking part in the retest. In this case, we informed participants that "Thank you for your contributions. If you want to participate in the test again, please write your name (you can use a nickname) and email address. We will send you a request again to participate in the study after 2-3 weeks. At this stage, you can give up the study at any time. Your personal information will be destroyed after the completion of the retest stage for confidentiality and safety purposes."

### Data Analysis

The data collected was exported from the online platform into the Statistical Package for the Social Science (SPSS) version 22.0 and Analysis of Moment Structures (AMOS) 25.0 for analysis. We considered a confidence interval of 95% ( $p < 0.05$ ).

For face validity, answers were evaluated descriptively. For content validity, Davis technique and the Content Validity Index (CVI) were used to evaluate the experts' opinions at item (I-CVI) and scale levels (S-CVI).<sup>24-27</sup> The number of experts rating each item as 3 or 4 was therefore divided by the total number of experts so that we could assess I-CVI. S-CVI was computed by summing the proportion of items that were rated 4 or 3 by the number of experts. The Kendall W analysis was used to test the level of agreement among experts.

Construct validity was assessed using Explanatory Factor Analysis (EFA) and Confirmatory Factor Analysis (CFA). The study sample was randomly split using participant entry codes; the first half (n= 300) was used to explore the measurement model with EFA, and the second half (n= 301) was used to confirm the model using CFA. Data suitability for factor analysis was examined using the Kaiser-Meyer-Olkin (KMO) coefficient and Barlett Sphericity test. We used Exploratory Principal Axis Factoring (PAF) with Oblimin rotation to explore the factor structure. Eigenvalues of 1 or higher and factor loadings of at least 0.30 were considered acceptable for factor and item retention, respectively.<sup>23,24,28</sup> For CFA, Pearson  $\chi^2$ , degree of freedom, root mean square error of approximation (RMSEA), goodness-of-fit index (GFI), comparative fit index (CFI), and normal fit index (NFI) were examined. Reliability of the final scale was determined using Cronbach's alpha, test-retest reliability (using Pearson's correlation, t-test, and intraclass correlation coefficient (ICC))<sup>23,29,30</sup>, item-total correlations, ceiling, and floor effects, and Hotelling's T-squared test for response bias.<sup>27,29</sup> To calculate the floor and ceiling effect the number of people who could get the floor/lowest (10/50) and highest/ceiling scores (50/50) on the instrument were summed up. It was calculated what percentage of

these made up the total sample. A minimum Cronbach's  $\alpha$  coefficient of  $>0.60$  was considered to indicate acceptable reliability.<sup>24,26</sup>

## Ethics

Written permission (via e-mail) was obtained from the first author of the original scale (Dr. Déborah Oliveira) to translate and validate the MOCHAD-10 for the Turkish population. Ethical approval was obtained from the Ethical Committee of Dokuz Eylul University (approval number: 2021/08-09 date: March 8, 2021). All participants were offered information about the study and gave their informed consent to take part. At the end of the survey, after answering all the study questions, participants were asked whether they wished to take part in the re-test phase within 2-3 weeks, for which we asked their names and email address. We informed them that they were still able to withdraw from the study at any time during that period and that their personal information would be destroyed after the completion of the retest stage for confidentiality and safety purposes. Only the research team had access to their data. Because participants provided their answers to the survey before giving any personal information (if they wished to), we believe their answers have not been affected by this.

## Results

### 1) Translation and back translation

After translation and back translation, nine of the ten items were nearly exactly similar to the original items and did not need to be altered. The item 7 was revised to better reflect the Turkish meaning (“When I think about dementia, I feel overwhelmed” was replaced with “When I think of dementia, I feel nauseous”) (Supplementary material).

### 2) Face and content validity

Face validity assessment was conducted with 15 individuals aged (mean)  $55.73 \pm 10.52$  years (range:40-70), 60% were female, 46.7% had a job, and over 73% reported their income status as being equal to expenditure. Participants' level of school attainment was similarly distributed among the education groups (n=3 primary school; n=3 high school; n=3 university degree; n=4 master's degree; n=2 doctorate). Participants reported taking 2-5 min to complete the T-MOCHAD-10. All participants stated that the number and order of questions were acceptable and did not have any problems with the rating of the statements. Most participants stated that the T-MOCHAD-10 was relevant and understandable. However, one participant raised the issue that some people may not know the meaning of the word dementia. To address this, we added to the first part of the survey a brief sentence referring to terms commonly used in the Turkish culture to refer to dementia (dementia is also known as...). In content validity assessment, the I-CVI for the 10 items was in the range of 0.85 and 1 and S-CVI was 0.98. No statistically significant differences were found between the scores given for each item (Kendall  $W = 0.09$ ,  $p = 0.60$ ). For

**Table 1** Socio-demographic profile of the respondents (n=601)

	n	%
<b>Sex</b>		
Male	208	34.6
Female	393	65.4
<b>Income</b>		
Income less than expenditure	86	14.3
Income equal to expenditure	321	53.4
Income more than expenditure	194	32.3
<b>Marital status</b>		
Married	502	83.5
Single	99	16.5
<b>Family member with dementia</b>		
Yes	70	11.6
No	531	88.4
<b>Caregiver of people with dementia</b>		
Yes	47	7.8
No	554	92.2
<b>Education level</b>		
Literate/elementary school	20	3.3
High school graduate	70	11.6
University degree	373	62.1
Postgraduate degree	138	23.0
<b>Mean age and SD</b>	53.86	9.31
<b>Age range of respondents</b>	40	85

this reason, all items were retained in the instrument at this stage.

### 3) Construct validation and reliability tests

The mean age of participants (n=601) was  $53.86 \pm 9.31$  years (range=40-85), 65.4% (n=393) were female, 83.5% (n=502) were married, 53.4% (n=321) reported their income status as being equal to expenditure, and over 11% (n=70) had a family member living with dementia (Table 1). We found a KMO coefficient of 0.79 and a Bartlett Sphericity test  $\chi^2$  of 959.58 ( $p < 0.001$ ), which demonstrated that the data were suitable for factor analysis. Within the EFA, two factors with Eigenvalues greater than 1 were determined: the first explained 29.97% and the second explained 14.74% of the total variance. Together, the two factors explained 44.71% of the total variance. Subsequent examination of the scree plot showed that a two-factor solution was optimal. Factor loadings ranged from 0.63 to 0.82 for factor 1 ('positive cues to action'), and from 0.37 to 0.70 for factor 2 ('negative cues to action') (Table 2).

The CFA applied to the two-factor solution initially suggested that the model was not appropriate. Both the CFI (=0.85) and the GFI (=0.87) had a lower than the acceptable value of 0.90 and 0.95, respectively. Errors related to covariances were addressed and the model then indicated a better fit. CFA revealed adequate factor loadings ranging from 0.42 to 1.50 (Figure 1). Model fit indicators were determined as follows: CFI=0.96, GFI=0.95, NFI=0.93, Chi-square/degree of freedom ( $\chi^2/df$ ) = 2.27,  $p < 0.001$ , and RMSEA=0.06 (Figure 1).



**Table 2** Factor loadings, Exploratory Principal Axis Factoring (PAF) with Oblimin rotation (n=300)


Items*	Factor 1:Positive cues to action	Factor 2:Negative cues to action	Domains from the Health Belief Model <sup>a</sup>
1. I am able to make differences that will change the risk of developing dementia	0.76		CA/SE
2. Changing my lifestyle and health habits can help me reduce my chance of developing dementia	0.82		CA/PB
3. Having risk factor(s) for dementia makes me think I have to change my lifestyle and behavior	0.68		CA
4. Learning more about dementia from the media makes me think I have to change my lifestyle and behavior	0.63		CA
5. Knowing family member(s) with dementia makes me think I have to change my lifestyle and behavior	0.65		CA
6. When I think about dementia my heart beats faster		0.70	CA/PSE
7. When I think about dementia I feel nauseous		0.67	CA/PSE
8. The thought of dementia scares me		0.62	CA/PSE
9. My feelings about myself would change if I develop dementia		0.37	CA/PSE
10. There is a strong possibility that I will develop dementia		0.43	CA/PSU
Explained variance (%)	29.97	14.74	

<sup>a</sup> SE: self-efficacy, PB: perceived benefits, CA: cues to action, PSE: perceived severity, PSU: perceived susceptibility

\* Turkish version of instrument was administered to the participants.

Using Pearson product-moment correlation, we found a moderate, positive correlation between positive cues to action and negative cues to action subscales, which was statistically significant ( $r=0.336$ ,  $p<0.001$ ) (Figure 2).

In the test-re-test (n=65), the mean score of the 65 participants was  $32.86\pm 5.57$  on the test and  $33.27\pm 5.84$  for the re-test. No significant differences were observed between the test and retest average scores ( $t = -0.98$ ,  $p=0.32$ ),



**Figure 1** Confirmatory Factor Analysis of T-MOCHAD-10




Figure 2 Correlations between the Factors

indicating the stability of measurements over time. A Pearson product-moment correlation was run to determine the relationship between total scores of T-MOCHAD-10 test and the re-test. There was a strong, positive correlation between the total scores of T-MOCHAD-10 test and the re-test, which was statistically significant ( $r=0.823$ ,  $p<0.001$ ) (Table 3). The T-MOCHAD-10 was also found to be reliable with respect to its test-retest reliability, with an intraclass correlation coefficient (ICC) of 0.903 (confidence interval [CI] 95%: 0.841-0.941) for the total scale, 0.763 (CI:0.611-0.856) for the positive cues to action subscale, and 0.873 (CI:0.793-0.923) for the negative cues to action subscale. The Cronbach’s  $\alpha$  score for the entire scale was 0.79 (‘positive cues to action’ = 0.82, ‘negative cues to action’ factor = 0.73). The Hotelling’s T-squared test was 4683.80, with a significance of  $p<0.001$ , demonstrating an absence of response bias. We also found no floor or ceiling effects

(=0.33%). All the item-total correlations were acceptable, ranging between 0.39 and 0.68: from 0.72 to 0.82 for positive cues to action and from 0.55 to 0.79 for negative cues to action (Table 4).

### Discussion

We aimed to develop and validate the Turkish version of the MOCHAD-10 scale for use in Turkey. Using face and content validity assessments, as well as exploratory and confirmatory factor analyses and reliability tests, we showed that the T-MOCHAD-10 generates valid and reliable scores of motivation to change lifestyle for dementia risk reduction among the Turkish population aged 40 and over. As with the original version, a 10-item two-factor structure produced the best psychometric performance.

This is the first study to translate and validate the MOCHAD-10 into another language and context, and the original study did not include face validity<sup>8</sup>, therefore we do not have any elements for comparison. However, our study demonstrated that the target population did not have difficulty in completing the scale, and that all the items were understandable and considered to be relevant. The content validity scores (I-CVI and S-CVI) were all above the minimally required<sup>30</sup>, which demonstrates that we were able to reach acceptable agreement among the experts and that the MOCHAD-10 appears to measure the concept it is intended to measure.

The two-factor structure of T-MOCHAD-10 was identical to the one found in the original study, demonstrating that the items had a high level of relationship with their respective factors and that both factors can adequately measure the construct they intend to measure in a strong factor structure, similarly to the original study.<sup>8</sup> After addressing some errors in

Table 3 Test-re-test of the T-MOCHAD-10 (n=65)

Variables	Test	Retest	t	p	r	p
Positive cues to action	19.55±3.69	19.52±3.55	0.07	0.93	0.779	<0.001
Negative cues to action	13.30±3.71	13.75±3.84	-1.42	0.15	0.614	<0.001
Total scores	32.86±5.57	33.27±5.84	-0.98	0.32	0.823	<0.001

t: paired sample t-test, r: Pearson product-moment correlation

Table 4 Item-total and item-subdimension correlation scores (n=601)

Subdimension	Item	Item–total correlation (r)*	Item–subdimension correlation (r)*
Positive cues to action	Item 1	0.58	0.77
	Item 2	0.62	0.82
	Item 3	0.68	0.79
	Item 4	0.61	0.72
	Item 5	0.61	0.71
Negative cues to action	Item 6	0.62	0.79
	Item 7	0.50	0.70
	Item 8	0.68	0.77
	Item 9	0.60	0.62
	Item 10	0.39	0.55

\*  $p < 0.001$

covariances, the CFA results indicated that the factor loadings and fit indices were all within the limits stated in the literature, confirming that the factor structure of the instrument was the best possible fit.<sup>31,32</sup> In general, the change in fit as a result of error correlation results in 0.02 to 0.03 betterment in fit across indices.<sup>33</sup> If correlated errors occur because of the researcher's aim to achieve a good model fit for their data, it means that the probability of correlated errors is reduced if a model already has an acceptable fit.<sup>34</sup> In this study, both the CFI (=0.85) and the GFI (=0.87) values were very close to the acceptable values of 0.90 and 0.95, respectively. For this reason, error covariances were addressed to indicate a better fit, with all fit indices being larger than the established threshold. The original version of MOCHAD-10 had similar fit indexes after addressing some errors in covariances (CFI=0.93, GFI=0.96, RMSEA=0.07,  $\chi^2/df=11.10$ ,  $p<0.001$ ).<sup>8</sup> These results indicate that the scale can provide valid outcomes of motivation to change lifestyle for dementia risk reduction among Turkish people aged 40 and above.

The T-MOCHAD-10 was found to provide highly reliable scores both by the test-re-test and by the internal consistency scores, meaning that the T-MOCHAD-10 can provide reliable outcomes when measuring the participants' motivation to change lifestyle for reducing the risk of dementia. The Cronbach's  $\alpha$  was similar to what was found in the original study (total  $\alpha=0.78$ , positive cues to action=0.81 and negative cues to action=0.70).<sup>8</sup> The Hotelling T-square test also indicated no significant response bias, suggesting that the participants answered questions based on their opinions, rather than on external factors.<sup>27,35</sup> We found a floor and ceiling effect of 0.33%, which is much under the limit of 20%, demonstrating an absence of such bias.<sup>27,35</sup>

## Conclusion

This study evaluated the psychometric properties of the Turkish version of the Motivation to Change Lifestyle for Dementia Risk Reduction (T-MOCHAD-10). The results obtained from this study showed that the MOCHAD-10 was a robust two-factor structure tool that provides valid and reliable outcomes reflecting people's motivation to change lifestyles for dementia risk reduction. By using this scale, health professionals will be able to assess people's motivation to change lifestyles for developing risk of dementia and engage in appropriate initiatives to enable people to implement these changes. Since this study involved a non-probabilistic sample and was conducted online, people without internet access were arguably not able to participate in the study. These may limit the wide generalization of the study results. Further studies should consider collecting data face-to-face to evaluate whether the participants understand the items and to allow for participation from a wider group of individuals.

## Acronyms

CFA: Confirmatory Factor Analysis; CFI: Comparative Fit Index; EFA: Exploratory Factor Analysis; GFI: Goodness-of-Fit Index; KMO: Kaiser-Meyer-Olkin; MOCHAD-10: Motivation to Change Behavior for Dementia Risk Reduction Scale, PAF:

Principal Axis Factoring, RMSEA: Root Mean Square Error of Approximation; T-MOCHAD-10: Turkish Version of Motivation to Change Behavior for Dementia Risk Reduction Scale.

## Author Statements

### Ethical Approval

Ethical approval was obtained from the Ethical Committee of Dokuz Eylül University (approval number: 2021/08-09 date: March 8, 2021).

### Funding

This study was not funded.

### Competing Interests

The authors declare no conflicts of interest.

### Acknowledgment

The authors are grateful to the participants for their participation in this study.

### Author Contributions

All authors were involved in the conception and design of the study. M.A.A., Ö.K., and B.A.S. performed the statistical analyses. M.A.A. wrote the first draft of the manuscript. All authors critically revised the manuscript for important intellectual content and approved the final version submitted.

### Data Availability Statement

All data generated or analyzed during this study are included in this article and its supplementary material files. Further inquiries can be directed to the corresponding author.

### Competing Interests

The authors declare no conflicts of interest.

### Acknowledgement

The authors are grateful to the participants for their participation in this study.

### Funding

This study was not funded.

## References

1. Prince MJ, Wu F, Guo Y, Robledo LMG, O'Donnell M, Sullivan R, et al. The burden of disease in older people and implications for health policy and practice. *The Lancet*. 2015;385(9967):549–62.

2. World Health Organization. Global action plan on the public health response to dementia 2017-2025. [accessed 21 October 2021]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/259615/9789241513487-eng.pdf?sequence=1>.
3. GBD 2019 Dementia Forecasting Collaborators. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: An analysis for the Global Burden of Disease Study 2019. *The Lancet Public Health*. 2022;7:2105–25.
4. World Health Organization. Risk reduction of cognitive decline and dementia, 2019. [accessed 21 October 2021]. Available from: [https://www.who.int/mental\\_health/neurology/dementia/guidelines\\_risk\\_reduction/en/](https://www.who.int/mental_health/neurology/dementia/guidelines_risk_reduction/en/).
5. Kim S, Sargent-Cox K, Cherbuin N, Anstey KJ. Development of the motivation to change lifestyle and health behaviors for dementia risk reduction scale. *Dement Geriatr Cogn Disord Extra*. 2014;4:172–83.
6. Abraham C, Sheeran P, et al. The health belief model. In: Ayers S, Baum A, McManus C, eds. *Cambridge Handbook of Psychology, Health and Medicine*, 2nd ed., Cambridge United Kingdom: Cambridge University Press; 2007:97–132.
7. Kim S, Sargent-Cox KA, Anstey KJ. A qualitative study of older and middle-aged adults' perception and attitudes towards dementia and dementia risk reduction. *J Adv Nurs*. 2015;71(7):1694–703.
8. Oliveira D, Aubeeluck A, Stupple E, Kim S, Orrell M. Factor and reliability analysis of a brief scale to measure motivation to change lifestyle for dementia risk reduction in the UK: the MOCHAD-10. *Health Qual Life Outcomes*. 2019;17:75.
9. Stretcher VJ, Rosenstock JM. The health belief model. *Health behavior and health education*. San Francisco: Jossey-Bass; 1997. p. 41–59.
10. Akyol MA, Zehirlioglu L, Erunal M, Mert H, Hatipoglu NS, Küçük-güçlü Ö. Determining middle-aged and older adults' health beliefs to change lifestyle and health behavior for dementia risk reduction. *Am J Alzheimers Dis Other Dement*. 2020;35:1–7.
11. Bosco A, Jones KA, Lorito CD, Stephan BC, Orrell M, Oliveira D. Changing lifestyle for dementia risk reduction: Inductive content analysis of a national UK survey. *Plos One*. 2020;15(12):e0244399.
12. Lee W, Gray SL, Zaslavsky O, Barthold D, Marcum ZA. Association between having a family member with dementia and perceptions of dementia preventability. *Aging Ment Health*. 2022;26(2):270–6.
13. Choi WH, Seo YM. Determinants of intention toward dementia-preventive behaviors among middle-aged Korean adults. *Int J Aging Hum Dev*. 2021. <https://doi.org/10.1177/00914150211024180>.
14. Kessler E-M, Bowen CE, Baer M, Froelich L, Wahl H-W. Dementia worry: A psychological examination of an unexplored phenomenon. *Eur J Ageing*. 2012;9:275–84.
15. Livia B, Elisa R, Claudia R, Roberto P, Cristina A, Emilia ST, et al. Stage of change and motivation to a healthier lifestyle before and after an intensive lifestyle intervention. *Journal of Obesity*. 2016. Article ID 6421265:1-7.
16. Becker MH. The Health Belief Model and personal health behavior. *Health Education Monographs*. 1974;2:324–508.
17. Rosenstock IM. Historical origins of the health belief model. *Health Education Monographs*. 1974;2:328–35.
18. Terwee CB, Bot SDM, de Boer MR, van der Windt, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol*. 2007;60:34–42.
19. Maneesriwongul W, Dixon JK. Instrument translation process: A methods review. *J Adv Nurs*. 2004;48(2):175–86.
20. Holden RR. Face validity. In: Weiner IB, Craighead WE, eds. *The Corsini encyclopedia of psychology*, Hoboken, NJ: Wiley; 2010:637–8.
21. Rouquette A, Falissard B. Sample size requirements for the internal validation of psychiatric scales. *Int J Methods Psychiatr Res*. 2011;20(4):235–49.
22. Polit DF. Getting serious about test–retest reliability: A critique of retest research and some recommendations. *Quality of Life Research*. 2014;23(6):1713–20.
23. Souza ACD, Alexandre NMC, Guirardello EDB. Psychometric properties in instruments evaluation of reliability and validity. *Epidemiol Serv Saude*. 2017;26:649–59.
24. Cam MO, Baysan-Arabaci L. Qualitative and quantitative steps on attitude scale construction. *Turkish J Res Dev Nurs*. 2010;12(2):59–71.
25. Crestani AH, Moraes AB, Souza APR. Content validation: Clarity/relevance, reliability and internal consistency of enunciative signs of language acquisition. *CoDAS*. 2017;29(4):e20160180.
26. Leech NL, Barrett KC, Morgan GA. *SPSS for intermediate statistics: Use and interpretation*. New York: L. Erlbaum Associates; 2008.
27. Nunnally JC, Bernstein IH. *Psychometric Theory*. 3rd ed. New York: McGraw-Hill; 2010.
28. Polit DF, Beck CT, Owen SV. Is the CVI an acceptable indicator of content validity? Appraisal and recommendations. *Res Nurs Health*. 2007;30(4):459–67.
29. Streiner DL, Kottner J. Recommendations for reporting the results of studies of instrument and scale development and testing. *J Adv Nurs*. 2014;70(9):1970–9.
30. Sousa VD, Rojjanasrirat W. Translation, adaptation and validation of instruments or scales for use in cross-cultural health care research: A clear and user-friendly guideline. *J Eval Clin Pract*. 2011;17(2):268–74.
31. DeVellis RF. *Scale Development, Theory and Applications*. 4th ed. New Delhi, India: SAGE Publication Inc; 2016.
32. Johnson B, Christensen L. *Educational Research: Quantitative, Qualitative, and Mixed Approaches*. p 199, 3rd ed. California: SAGE Publication; 2014. p. 199–222.
33. Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Modeling*. 1999;6(1):1–55.
34. Hermida R. The problem of allowing correlated errors in structural equation modeling: Concerns and considerations. *Comput Methods Soc Sci*. 2015;3(1):5–17.
35. Rattray J, Jones MC. Essential elements of questionnaire design and development. *J Clin Nurs*. 2007;16:234–43.