





RESEARCH
ARTICLE

-  Seher Palanbek Yavas¹
 Caner Baysan²
 Ayse Emel Onal³
 Ozkan Ayvaz³

¹ Çanakkale Onsekiz Mart
University, Faculty of Medicine,
Department of Public Health,
Çanakkale, Türkiye

² Izmir Democracy University,
Faculty of Medicine, Department of
Public Health, Izmir, Türkiye

³ Istanbul University, Faculty of
Medicine, Department of Public
Health, İstanbul, Türkiye

Corresponding Author:

Seher Palanbek Yavas

mail: seher.palanbekyavas@istanbul.edu.tr

Received: 11.05.2022

Acceptance: 26.09.2022

DOI: 10.18521/ktd.1113580

Konuralp Medical Journal

e-ISSN1309-3878

konuralptipdergi@duzce.edu.tr

konuralptipdergisi@gmail.com

www.konuralptipdergi.duzce.edu.tr

Turkish Adaptation of the Mental Health Literacy Scale for Healthcare Students: A Study of Validity and Reliability**ABSTRACT**

Objective: This study has aimed to investigate the validity and reliability of the Turkish version of the Mental Health Literacy in Healthcare Students (MHLS-HS).

Methods: The study sample comprises 275 students at Istanbul Medical Faculty between the ages of 18 and 27. Data were collected using the MHLS-HS, the Beliefs about Mental Illness Scale (BMI), and the Positive Mental Health Scale (PMS). For language adaptation, translation, back translation, expert comments, and a trial application were conducted. Validity was determined using the content validity index and confirmatory factor analysis, while reliability was determined using Cronbach's alpha analysis and the test-retest method.

Results: The content validity index of the scale was found to be 0.99 and the item loads were found to be 0.38-0.94 according to confirmatory factor analysis. The Cronbach's alpha coefficient was found to be 0.79 for the whole scale and between 0.60-0.89 for the five sub-dimensions. In equivalent criterion validity, it was determined that BMI had a weak negative correlation ($r=-0.360$, $p<0.001$) and a weak positive correlation ($r=0.327$, $p<0.001$) with PMS. A statistically significant difference was found in the sub-dimensions of Maintaining Positive Mental Health, Stigmatizing Mental Illnesses, and Seeking Help Effectiveness between those with and without mental illness in the discrimination analysis of the RSS-SS scale ($p<0.001$, $p=0.029$, $p=0.034$, respectively).

Conclusions: The Mental Health Literacy Scale in Healthcare Students was found to be valid and reliable. It can be used to assess and improve the educational experience of medical school students.

Keywords: Mental Health Literacy, Validity, Reliability, Healthcare Students, Scale.

Sağlık Öğrencileri İçin Ruh Sağlığı Okuryazarlık Ölçeğinin Türkçeye Uyarlanması: Geçerlik ve Güvenirlik Çalışması**ÖZET**

Amaç: Bu çalışmada, Sağlık Öğrencileri için Ruh Sağlığı Okuryazarlık Ölçeğinin (RSOÖ-SÖ) Türkçe uyarlanması için geçerlilik ve güvenilirlik yapılması amaçlanmıştır.

Gereç ve Yöntem: Çalışma örneklemini 18-27 yaş arasında İstanbul Tıp Fakültesinde öğrenim gören 275 kişiden oluşmaktadır. RSOÖ-SÖ ölçeği, Ruhsal Hastalıklara Yönelik İnanç Ölçeği (RHYİÖ) ve Pozitif Mental Sağlık Ölçeğini (PMSÖ) veri toplamada kullanılmıştır. Dil uyarlaması için çeviri, geri çeviri, uzman görüşleri ve pilot uygulama yapılmıştır. Geçerlik için kapsam geçerlilik indeksi, doğrulayıcı faktör analizi ve güvenilirliği test etmek için Cronbach alfa analizi, test-tekrar test yöntemi kullanılmıştır.

Bulgular: Ölçeğin kapsam geçerlilik indeksi 0.99 ve doğrulayıcı faktör analizine göre madde yükleri 0.38-0.94 olduğu bulunmuştur. Cronbach alfa katsayısı tüm ölçek için 0.79, beş alt boyutu için ise 0.60-0.89 arasında olduğu belirlenmiştir. Eşdeğer ölçüt geçerliliğinde RHYİÖ ile negatif yönlü zayıf korelasyon ($r=-0.360$, $p<0.001$), PMSÖ ile ise pozitif yönlü zayıf korelasyon ($r=0.327$, $p<0.001$) gösterdiği saptanmıştır. RSOÖ-SÖ ölçeğinin ayırt edicilik analizinde ruhsal hastalığı olan ve olmayanlar arasında Pozitif Ruh Sağlığını Sürdürme, Ruhsal Hastalıkları Damgalama ve Yardım arama etkinliği alt boyutlarında istatistiksel olarak anlamlı fark saptanmıştır (Sırasıyla $p<0.001$, $p=0.029$, $p=0.034$).

Sonuç: Sağlık Öğrencileri için Ruh Sağlığı Okuryazarlığı Ölçeğinin geçerli ve güvenilir olduğu saptanmıştır. Tıp fakültesi öğrencilerinde eğitimin değerlendirilmesi ve iyileştirilmesi için kullanılabilir.

Anahtar Kelimeler: Ruh Sağlığı Okuryazarlığı, Geçerlik, Güvenirlik, Sağlık Öğrencileri, Ölçek.

INTRODUCTION

The World Health Organization (WHO) describes mental health as "a condition of well-being that an individual recognizes his or her strengths, can handle the usual demands of life, might work effectively, and also can make a significant contribution to his or her society"(1). Worldwide, an approximated 322 million individuals suffer from depression, 264 million from anxiety, 45 million from bipolar illness, 20 million from schizophrenia, and 50 million from dementia (2,3). More than 70% of these people are unable to get help from health professionals as they avoid and delay treatment due to a lack of knowledge to define mental illness, being unaware of how to access treatment, prejudice towards those with mental illness and fear of stigmatization (4,5).

Mental health literacy is a significant factor in mental health and has the order to enhance public mental health. (6). Studies have reported that knowing mental health and diseases increases one's awareness in seeking help and treatment, benefiting from health services and early diagnosis and reducing mental illness stigma in society (7,8) Several scales such as the Vignette Interview, the Mental Health Literacy Scale (MHLS), and the Mental Health Literacy Scale for Youth and Young Adults (MHLq) have been developed to determine one's mental health literacy (9-12). These scales were created to measure and examine the literacy levels of the public, but various studies have shown that mental health literacy should also be improved in healthcare professionals and students (medical school students, nursing students, health vocational school students, etc.) (13). Healthcare professionals and students have high levels of stress, burnout, anxiety and depression (14,15). Healthcare professionals and students exhibit a pervasive stigma, as well as unfavourable attitudes and beliefs, toward those with mental illness. Therefore, healthcare professionals and students may refuse to explain and ignore their mental problems and avoid seeking help and treatment (16,17). In addition, studies have reported that mental illnesses can be overlooked in primary care centres, whereby their diagnosis rates are low (18,19).

Previous studies have suggested identifying and improving mental health literacy among healthcare professionals and students. However, there is no Turkish scale to determine mental health literacy for healthcare students. Mental Health Literacy Scale for Healthcare Students, which covers five essential components of health literacy (maintaining healthy mental health, recognizing mental illness, a stigma attitude, help-seeking efficacy, and help-seeking attitude), may fill this gap (13). This study aimed to adapt the Mental Health Literacy Scale for Healthcare Students into Turkish and to determine its Turkish validity and reliability.

MATERIAL AND METHODS

Study Design and Sample Size: This methodological study was undertaken with healthcare students aged 18 to 27 who studied at the Istanbul University Faculty of Medicine during the educational year 2021-2022 and volunteered to participate. The data were collected online between November 2021 and March 2022. Filling out the questionnaire lasted around 20 minutes. The sample size in scale adaptation studies should be at least 5-10 times the number of scale items. Therefore, a total of 275 people were reached in this study (the number of scale items = 26) (20). Two weeks after the first phase of data collection, the questionnaire was sent again for re-testing to health students who already filled out it, and a total of 104 healthcare students responded. Ethical approval was received by the Social and Humane Ethics Committee (Dated: 30/03/2021 E-155284) at Istanbul University, and permission from Yin-Ju Lien, the co-author of the article, via e-mail, to use and verify the validity and reliability of the scale in Turkish (February 26, 2021).

Data Collection Tools: The data were gathered using a personal information form that included questions on healthcare students' sociodemographic characteristics of healthcare students and the Mental Health Literacy Scale for Healthcare Students (MHLS-HS). The data collection form included two additional scales, namely Positive Mental Health Scale (PMSÖ) and the Beliefs toward Mental Illness Scale (RHYİÖ), to examine the equivalent criterion validity of the scale.

The Mental Health Literacy Scale for Healthcare Students (MHLS-HS): The MHLS-HS was designed by Chao et al. utilizing a total of 1294 medical and public health undergraduate students from 11 universities to assess mental health literacy among healthcare students (13). It consists of 26 items and five 5 subscales (M = Maintenance of positive mental health 10 items; R = Recognition of mental illness 4 items; S = Mental illness stigma attitude 6 items; HE = Help-seeking efficacy 3 items; HA = Help-seeking attitude 3 items). This is a 5-point Likert type scale, scoring from 1 (strongly disagree) to 5 (strongly agree). The scale is interpreted through total and subscale scores, where the subscale of "S= Mental illness stigma attitude" is scored in reverse. A higher score indicates better mental health literacy. The MHLS-HS has good reliability. The Cronbach's alpha coefficient for the whole scale was 0.81 and varied from 0.70 to 0.87 for its subscales. The correlation of the MHL-HS with the Social Distance Scale and the Positive Mental Health Scale, which were used to achieve similar scale validity, supported the convergent validity of the MHL-HS, and higher MHL-HS total scores were significantly associated with lower social distance toward people with

mental illness ($r = -0.26$, respectively; $p < 0.001$, $r = 0.35$; $p < 0.001$) (13).

Positive Mental Health Scale (PMSÖ):

Positive Mental Health Scale (PMHS) was developed by Lukat et al. (2016) and adapted into Turkish by Yılmaz-Akbaba and Eldeleklioğlu (2019) (21,22). It consists of nine items, scoring from "Not True (1) to "True (4)". The internal consistency coefficient of the PMSÖ was determined to be 0.85, and substantial positive connections were established between the PMSÖ and the scales used for similar scale validity.

Beliefs toward Mental Illness Scale (RHYİÖ): The Beliefs towards Mental Illness Scale (BMI) was created by Hirai and Clum (2000) and adapted into Turkish by Bilge and Çam (2008) (23,24). This is a 6-point Likert-type scale, scoring from "Totally Disagree (0)" to "Totally Agree (5)". It consists of 21 items and three subscales, involving danger, lack of social and interpersonal skills, and incurability. The Cronbach's alpha coefficient was 0.82 for the entire scale and ranged from 0.69 to 0.80 for the subscales.

Translation of the Scale and a Pilot Application: The steps of translation, back-translation and pilot application were followed sequentially to create the Turkish form of the scale. A professional translator who graduated in English language literature (C.B.) translated the scale from English to Turkish. A foreign citizen whose mother tongue is English and who speaks Turkish (K.Y.) translated it back from Turkish to English.

Upon the consensus of the researchers, the scale items were evaluated and scored by a total of seven experts, including six physicians and one psychiatric nurse, in terms of their accuracy, suitability and intelligibility for health students (M.Ç., A.M., L.B.A., S.U.U., E.Ç.K., Ü.Z., G.A.M.). Davis method was used to assess items on the scale, scoring "inappropriate (1)", "should be seriously reviewed (2)", "should be slightly revised (3)", and "appropriate (4)" (25). The content validity ratio (CVR) of the scale items ranged from 0.86 to 1.00, and the content validity index (CGI) was calculated to be 0.99.

A pilot application was conducted with a group of 30 healthcare students to evaluate the intelligibility of the scale. The data obtained from the pilot application were not used in the study.

Data Analysis: Before the analysis, the presence of missing data in the data set was checked (even though the scale was set in a way that respondents could not answer the next question without answering the previous one, thus the possibility of missing data was eliminated). The Kolmogorov-Smirnov test was utilized to determine whether or not the data has a normal distribution. The distribution of the data was not normal. As the data lacked a normal distribution, the Mann Whitney U test and Spearman's correlation analysis were applied for statistical analysis. The internal

consistency and reliability of the scale and its subscales were evaluated using Cronbach's alpha. Correlation analysis and Interclass Correlation (ICC) analysis were used to compare the test-retest scale scores for time invariance. The construct validity of the scale was determined using confirmatory factor analysis (CFA). Davis method was used to calculate the CGI and CVR values, where the limit values for CVR and CGI should be above 0.80 and 0.67, respectively (25). PMSÖ and RHYİÖ were used for the convergent and discriminant validity of the scale. A p-value less than 0.05 was considered statistically significant. On the scale, items with reverse scores were scored in reverse. The IBM SPSS (Social Sciences Statistical Package) 20 package program, AMOS and Lisrel program were used to examine the data.

RESULTS

Characteristics of the Participants: The mean age of 275 healthcare students who agreed to participate in the study was 20.9 ± 2.1 years (range, 18-27), 59.3% of them were women. In addition, 26.5% of them ($n=73$) were 1st graders, 19.6% ($n=54$) 2nd graders, 12.4% ($n=34$) 3rd graders, 7.6% ($n=21$) were 4th graders, 16.4% ($n=45$) were 5th graders and 17.5% ($n=48$) were 6th graders, 9.5% ($n=26$) they had chronic health issues, and 14.9% ($n=41$) had mental illness in the past.

Validity Analysis of the Mental Health Literacy Scale for Healthcare Students (RSOÖ-SÖ):

Content Validity: The scale was evaluated for content validity by seven field experts (M.Ç., A.M., L.B.A., S.U.U., E.Ç.K., Ü.Z., G.A.M.). The content validity ratio (CVR) of the scale items ranged between 0.86 and 1.00, and the content validity index (CGI) of the scale was determined as 0.99.

Construct Validity: A confirmatory factor analysis (CFA) was performed for the construct validity of the RSOÖ-SÖ. As a result, an item (numbered 4) in the subscale of "Maintenance of Positive Mental Health (M)", stating "*having religious or spiritual beliefs*", had a factor loading as 0.07, and an item (numbered 3) in the subscale of "Recognition of Mental Illness (R)", stating "*if one needs higher doses of a drug to achieve the same effect; to what extent do you think he/she is likely to have a substance-related addiction?*", had factor loading as 0.20. These two items were excluded from the scale due to low factor loadings. The CFA was conducted over 24 items and 5 subscales. CFA fit indices of the scale were as follows: χ^2/df 1.46, Root mean square error of approximation (RMSEA) = 0.041, Comparative fit index (CFI) = 0.96, Incremental fit index (IFI) = 0.96, Non-normed fit index (NNFI) = 0.95, Goodness-of-fit index (GFI) = 0.90, Standardized root mean square residual (SRMR) = 0.056, and Critical N (CN) = 219.37. Factor loadings of the scale items ranged between 0.38 and 0.94. Figure 1 shows the confirmatory factor analysis of the scale.




Figure 1. Confirmatory factor analysis diagram of the Mental Health Literacy scale for Healthcare Students

Equivalence Validity of the RSOÖ-SÖ:
 For the equivalence validity of the Mental Health Literacy Scale for Healthcare Students, its correlation with RHYİÖ and PMSÖ was evaluated, whereby the RSOÖ-SÖ had a weak negative correlation with RHYİÖ ($r=-0.360$, $p<0.001$) and a

weak positive correlation with PMSÖ ($r=0.327$, $p<0.001$) (Table 1).

Differential Validity of the RSOÖ-SÖ:
 Table 2 compares the RSOÖ-SÖ and subscales scores of healthcare students according to the presence of mental illness.

Table 1. The correlation coefficients of the Beliefs towards Mental Illness Scale and the Positive Mental Health Scale of the RSOÖ-SÖ Scale and its sub-dimensions

	RHYİÖ	PMSÖ
MHLS-HS	-0.360**	0.327**
M	-0.028	0.125
R	-0.102	0.395**
S	-0.620**	-0.404**
HE	-0.114	0.651**
HA	-0.200**	0.523**

M = Maintenance of positive mental health R = Recognition of mental illness; Mental illness stigma attitude; HE = Help-seeking efficacy; HA = Help-seeking attitude; RHYİÖ= Beliefs towards Mental Illness Scale; PMSÖ = Positive Mental Health Scale ** p<0.001.

Table 2. Comparison of the RSOÖ-SÖ Scale and sub-dimension scores of the students according to the presence of mental illness

	Mental illness		p
	None (n=234)	Yes (n=41)	
M, median (1Q-3Q)	33(29-36)	28(24-32)	<0.001
R, median (1Q-3Q)	12(11-14)	13(12-14)	0.371
S, median (1Q-3Q)	21(19-24)	23(20-26.5)	0.029
HE, median (1Q-3Q)	12(9-14)	12(10.5-15)	0.034
HA, median (1Q-3Q)	10(8-12)	12(9-14)	0.070
MHLS-HS, mean±sd	88(80-94.3)	88(79-94)	0.837

M = Maintenance of positive mental health R = Recognition of mental illness; Mental illness stigma attitude; HE = Help-seeking efficacy; HA = Help-seeking attitude

Reliability Analysis of the RSOÖ-SÖ:
Internal Consistency Analysis of the RSOÖ-SÖ: The Cronbach's alpha reliability coefficient was found to be 0.796 for the total scale and ranged between 0.608 and 0.899 for its five

subscales (Table 3). The item-total correlations were weak, moderate and positive (r=0.241-0.517). The correlation between the total scale and subscales scores ranged between 0.348 and 0.676 (p<0.001) (Table 4).

Table 3. The item score averages, factor loadings and internal consistency reliability

Sub-Dimensions	Items	Median (1Q-3Q)	Factor Loadings	Cronbach's alpha
M	M1	4(3-4)	0.59	0.826
	M2	4(3-4)	0.62	
	M3	4(3-4)	0.55	
	M5	4(3-5)	0.64	
	M6	4(2-5)	0.52	
	M7	4(3-4)	0.53	
	M8	4(3-4)	0.54	
	M9	3(2-4)	0.62	
	M10	3(2-4)	0.69	
	R	R1	3(3-4)	
R2		4(4-5)	0.67	
R4		4(4-5)	0.42	
S1		4(3-5)	0.38	
S	S2	1(1-2)	0.79	0.810
	S3	2(1-3)	0.55	
	S4	3(3-4)	0.80	
	S5	2(2-3)	0.61	
	S6	3(2-4)	0.75	
	HE1	2(1-2)	0.84	
HE	HE2	4(3-4)	0.94	
	HE3	4(3-5)	0.83	
	HA	HA1	4(3-5)	0.65
HA2		4(3-5)	0.92	
HA3		3(3-4)	0.61	
RSOÖ-SÖ score			88(80-94)	0.796

Table 4. Correlation coefficients of the scores of the RSOÖ-SÖ Scale and its sub-dimensions

Sub- dimensions	M	R	S	HE	HA
R	0.081	-	-	-	-
S	0.043	0.130*	-	-	-
HE	0.114	0.233**	0.085	-	-
HA	0.088	0.226**	0.211**	0.326**	-
MHLS-HS	0.676**	0.348**	0.504**	0.532**	0.548**

M = Maintenance of positive mental health R = Recognition of mental illness; Mental illness stigma attitude; HE = Help-seeking efficacy; HA = Help-seeking attitude; p<0.05, ** p<0.001.

Test-Retest Reliability of the RSOÖ-SÖ:

Two weeks later, the scale was reapplied to a total of 104 participants (50.5% (n=52) male) and the intragroup correlation coefficient (ICC) was calculated. The ICC was 0.894 for the total scale score (CI: 0.844-0.929). The total score correlation was found to be positive and very strong (r=0.875, p<0.001).

DISCUSSION

Mental health literacy is crucial to understanding how to improve and sustain positive mental health, identifying mental illnesses and treatment options, minimizing the drip of various psychiatric, increasing help-seeking efficacy and developing self-management skills, and is associated with the events frequently encountered in the course of a lifetime. This study examined the validity and reliability of the Mental Health Literacy Scale for Healthcare Students (MHLS-HS) in Turkish culture.

Discussion of Validity Analysis: A confirmatory factor analysis, which is an effective tool in assessing whether the factor model in the original scale is compatible with the data of the study was conducted to measure the construct validity of the scale (26). As a result, an item (numbered 4) in the subscale of "Maintenance of Positive Mental Health (M)" and an item (numbered 3) in the subscale of "Recognition of Mental Illness (R)" was excluded from the scale due to low factor loadings (0.07 and 0.20, respectively). Therefore, the subscale of Maintenance of Positive Mental Health included nine items, the subscale of Recognition of Mental Illness included three items, and the number of items in other subscales did not change. The factor loadings of the scale's last 24 items varied from 0.38 to 0.94. The factor loadings of scale items may vary by study. The factor loadings of the items in the scale were determined as 0.45 and above by Büyüköztürk et al. (27) and between 0.32-0.44 by Dede and Yaman (28). In our research, just two scale items had factor loadings of 0.38 and 0.42, while the remaining items had acceptable, very good, and excellent factor loading fit. Chao et al. found that the factor loadings ranged from 0.41 to 0.95(13). The CFA analysis revealed $\chi^2/df = 1.46$, RMSEA=0.041, IFI=0.96, CFI=0.96, GFI= 0.90, and SRMR=0.056, where $\chi^2/df \leq 3$, RMSEA \leq 0.05, IFI \geq 0.95 and GFI \geq 0.90 indicated good fit of the scale and CFI \geq 0.95 and SRMR \leq 0.08

indicated acceptable fit of the scale (29). Therefore, the CFA and goodness-of-fit statistics of the scale items in our study were good and at the desired level.

For equivalent criterion validity of the scale, its correlation with RHYİÖ was analyzed and a weak negative correlation was identified between them (r=-0.360, p<0.001). A high score on the RHYİÖ indicates a negative belief (24). Therefore, we can conclude that as the mental health literacy of health students increases, their negative attitudes towards mental illness decrease. In addition, for the equivalent criterion validity of the scale, its correlation with PMSÖ was also evaluated and a weak positive correlation was identified between them (r=0.327, p<0.001). A high PMSÖ score suggests great positive mental health. (22). Help-seeking efficacy and help-seeking attitude, two subscales of the RSOÖ-SÖ, showed higher correlations with the PMSÖ (0.651 p<0.001 and 0.523 p<0.001, respectively). Chao et al. (13) found a low positive correlation between the scale and the PMSÖ (r=0.35 p < 0.001).

Discussion of Reliability Analysis: The item-total score correlations of 24 items were positive, weak and moderate (r=0.241-0.517). The correlation coefficient between the overall scale and the five subscales ranged from 0.348 to 0.676. Positively significant relationships between the whole scale and subscale scores were obtained (p<0.001). This consequence theoretically coincides with the fact that as the mental health literacy level of healthcare students increases, they are willing to expand their knowledge about achieving and sustaining excellent mental health, their awareness of mental disorders and treatments, and their attitudes toward stigma reduction and help-seeking effectiveness.

The overall scale's Cronbach's alpha reliability coefficient (α) was determined to be 0.79 and ranged between 0.60 and 0.89 for its subscales, where $0.60 \leq \alpha \leq 0.80$ indicates that the scale is quite reliable (30). According to Chao et al., Cronbach's alpha coefficient was 0.81 for the entire scale and 0.70-0.87 for its subscales (13). Accordingly, our results comply with those reported by Chao et al.(14). In the adaptation study of the Mental Health Literacy Scale (MHL) to young adults (MHL-q), Cronbach's alpha was found to be 0.84 for the overall scale and 0.60 to 0.74 for the subscales. However, the items (29 items) and subscales (4

subscales) of the MHL-q are quite different from those in our scale (12).

The scale was re-applied two weeks later to assess its time consistency, and its first and second application scores were found to have a strong and positive correlation ($r=0.875$, $p<0.001$). An ICC analysis was performed to evaluate the agreement between test and retest scores of the scale, which was determined to have statistical significance ($p<0.001$). Thus, the scale was found to have test-retest reliability.

Discussion on the Relationship between Descriptive Characteristics and Subscales of the Scale: The healthcare students with mental illness had lower scores on the subscales of maintaining psychological wellbeing, mental disease stigma attitude, and higher scores on the subscale of help-seeking efficacy than those without mental illness, and the difference between them was statistically significant (respectively <0.001 , 0.029 , 0.034). Studies have indicated that persons with greater mental health literacy have better mental health and a less social distance from those with mental illness (30,31).

Items on the subscale of Maintenance Positive Mental Health subscale (M) of the RSOÖ-SÖ include items to positively affect competence, autonomy and mental health. Studies have reported that meeting these basic psychological needs leads to a better quality of life and mental health (32). Therefore, healthcare students, who will become health professionals in the future, should be provided with better mental health and better patient care (33).

The subscale of Recognition of Mental Illnesses (R) of the RSOÖ-SÖ includes items to identify mental illnesses such as schizophrenia, anxiety disorders, depression etc. Several mental disorders are common both in Turkey (34) and across the world (18). Increasing the knowledge of healthcare students on recognition of mental illness is important for early diagnosis and treatment when they graduate and start providing health services.

Another subscale of the RSOÖ-SÖ is the mental illness stigma attitude. Jorm and Kitchener have found that mental health literacy interventions and knowledge about mental illnesses reduce stigma (35). Link and Phelan explained that recognition of mental illness and labelling can have both positive and negative aspects, where labelling

other persons can be stigmatizing and labelling oneself can be a facilitator in treatment (36). If healthcare students can easily recognize mental illnesses and avoid labelling and stigmatization, they can accelerate patients' recovery. Discriminating tendencies of healthcare students can be detected using the scale and relevant educational interventions can be applied to them (13).

Strengths and Limitations: According to our knowledge, the RSO-S is the first instrument particularly created to measure mental health literacy among students in the healthcare field. Public healthcare students participated in the study of the original scale, but this study was conducted only with medical faculty students. However, there are no public healthcare students in undergraduate education in Turkey as the public health speciality is considered a separate branch after medical school. Therefore, we did not include public health assistants to avoid bias in the sample selection. In addition, our study does not represent all healthcare students in Turkey as it was conducted in a single faculty. Further studies can use different healthcare student groups and can be performed in multi-centred faculties. This is a self-report scale, therefore, even though invisibility and clandestinity are ensured during the study, the measurement of stigmatization towards mental health may not be completely independent of social desirability bias.

CONCLUSION

Our research suggests that the Turkish version of the RSOÖ-SÖ is a valid and reliable instrument with an adequate model fit. The RSOÖ-SÖ captured the multidimensionality appropriately, and the analysis confirmed the suitability of its five-factor structure. This scale encompasses both mental health promotion and many dimensions of mental health, including stigma and awareness of the mental illness. It allows rapid and comprehensive evaluation of mental health literacy among healthcare students and to conduct remedial curriculum studies in medical schools.

Acknowledgments: We appreciate the evaluation of Meltem Çöl, Aliye Mandracioğlu, Leyla Baysan Arabacı, Süleyman Utku Uzun, Ebru Çobanoğlu Karaca, Ülkü Zengin and Gül Anıl Mutlu at stage of scale construct validity of investigation.

REFERENCES

1. World Health Organization. Promoting mental health: concepts, emerging evidence, practice (Summary Report). [Internet]. Switzerland. World Health Organization; 2004. [cited 2022 March 20]. Available from: <https://apps.who.int/iris/handle/10665/42940>
2. World Health Organization. [Internet]. Switzerland. Depression and Other Common Mental Disorders: Global Health Estimates. World Health Organization; 2017. [cited 2022 March 20]. Available from: <https://apps.who.int/iris/handle/10665/254610>.
3. James SL, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 Diseases and Injuries for 195 countries and

- territories, 1990-2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392(10159):1789–858.
4. Thornicroft G. Most people with mental illness are not treated. *Lancet*. 2007;370(9590):807–8.
 5. Henderson C, Evans-Lacko S, Thornicroft G. Mental illness stigma, help-seeking, and public health programs. *Am J Public Health*. 2013;103(5):777–80.
 6. Kelly CM, Jorm AF, Wright A. Improving mental health literacy as a strategy to facilitate early intervention for mental disorders. 2007;187(7):1–5.
 7. Wei Y, McGrath PJ, Hayden J, Kutcher S. Mental health literacy measures evaluating knowledge, attitudes and help-seeking: A scoping review. *BMC Psychiatry* [Internet]. 2015;15(1):291
 8. Picco L, Abdin E, Pang S, Vaingankar JA, Jeyagurunathan A, Chong SA, et al. Association between recognition and help-seeking preferences and stigma towards people with mental illness. *Epidemiol Psychiatr Sci*. 2018;27(1):84–93.
 9. Jorm AF, Korten AE, Jacomb PA, Christensen H, Rodgers B, Pollitt P. “Mental health literacy”: A survey of the public’s ability to recognise mental disorders and their beliefs about the effectiveness of treatment. *Med J Aust*. 1997;166(4):182–6.
 10. O’Connor M, Casey L. The Mental Health Literacy Scale (MHLS): A new scale-based measure of mental health literacy. *Psychiatry Res*. 2015;229(1–2):511–6.
 11. Campos L, Dias P, Palha F, Duarte A, Veiga E. Development and psychometric properties of a new questionnaire for assessing mental health literacy in young people. *Univ Psychol*. 2016;15(2):61–72.
 12. Dias P, Campos L, Almeida H, Palha F. Mental health literacy in young adults: Adaptation and psychometric properties of the mental health literacy questionnaire. *Int J Environ Res Public Health*. 2018;15(7).
 13. Chao HJ, Lien YJ, Kao YC, Tasi IC, Lin HS, Lien YY. Mental Health Literacy in Healthcare Students: An Expansion of the Mental Health Literacy Scale. *Int J Environ Res Public Health*. 2020;17(3):1–15.
 14. Quek TTC, Tam WWS, Tran BX, Zhang M, Zhang Z, Ho CSH, et al. The global prevalence of anxiety among medical students: A meta-analysis. *Int J Environ Res Public Health*. 2019;16(15). :2735. DOI: 10.3390/ijerph16152735.
 15. Wilkes C, Lewis T, Brager N, Bulloch A, MacMaster F, Paget M, et al. Wellbeing and mental health amongst medical students in Canada. *Int Rev Psychiatry*. 2019;31(7–8):584–7.
 16. Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, et al. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet*. 2016;387(10023):1123–32.
 17. Lien YY, Lin HS, Lien YJ, Tsai CH, Wu TT, Li H, et al. Challenging mental illness stigma in healthcare professionals and students: a systematic review and network meta-analysis. *Psychol Heal*. 2021;36(6):669–84.
 18. Bor JS. Among the elderly, many mental illnesses go undiagnosed: Few health care providers have the training to address depression, anxiety, and other conditions in their older patients. *Health Aff*. 2015;34(5):727–31.
 19. Liu CH, Zhang E, Wong GTF, Hyun S, Hahm HC. Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: Clinical implications for U.S. young adult mental health. *Psychiatry Res*. 2020 Aug;290:113172.
 20. Petscher Y, Schatschneider C CD. *Applied quantitative analysis in education and the social sciences*. 1st Editio. New York: Routledge; 2013. p.169–242.
 21. Lukat J, Margraf J, Lutz R, Der Veld WM, Becker ES. Psychometric properties of the positive mental health scale (PMH-scale). *BMC Psychol*. 2016;4(1):1–14.
 22. Akbaba AY, Eldeleklioglu J. Adaptation of Positive Mental Health Scale into Turkish: A Validity and Reliability Study. *J Fam Couns Educ*. 2017;4(1):44–54.
 23. Hirai M, Clum GA. Development, Reliability, and Validity of the Beliefs Toward Mental Illness Scale. *J Psychopathol Behav Assess*. 2000;22:221–36.
 24. Bilge A, Çam O. Ruhsal Hastalığa Yönelik İnançlar Ölçeği’nin geçerliliği ve güvenilirliği. *Anadolu Psikiyatr Derg*. 2008;9(2):91–6.
 25. Reha A. Spor, Sağlık ve Eğitim Bilimlerinden Örneklerle Uygulamalı İstatistik ve Geçerlik-Güvenirlik. Ankara: Detay Yayınları; 2018.
 26. Floyd FJ, Widaman KF. Factor analysis in the development and refinement of clinical assessment instruments. *Psychol Assess*. 1995;7(3):286–95.
 27. Büyüköztürk Ş. *Data Analysis Handbook for Social Sciences, Statistics, Research Design SPSS Applications*. 23 th. Ankara: Academy Press; 2017.
 28. Dede Y, Yaman S. A Questionnaire for Motivation toward Science Learning : A Validity and Reliability Study. *Necatibey Faculty of Education, Electronic Journal of Science and Mathematics Education*. 2008;2(1):19–37.
 29. Kartal M, Bardakçı S. *SPSS ve AMOS uygulamalı örneklerle güvenirlik ve geçerlik analizleri*. Ankara: Akademisyen Kitabevi; 2018. p.186.

30. Jorm AF. Mental health literacy: empowering the community to take action for better mental health. *Am Psychol.* 2012;67(3):231–43.
31. Svensson B, Hansson L. How mental health literacy and experience of mental illness relate to stigmatizing attitudes and social distance towards people with depression or psychosis: A cross-sectional study. *Nord J Psychiatry.* 2016;70:309–13.
32. Ryan RM, Patrick H, Deci EL, Williams GC. Facilitating health behaviour change and its maintenance: Interventions based on self-determination theory. *Eur Heal Psychol.* 2008;10:2–5.
33. Murden F, Bailey D, Mackenzie F, Oeppen RS BP. The impact and effect of emotional resilience on performance: an overview for surgeons and other healthcare professionals. *Br J Oral Maxillofac Surg.* 2018;56(9):786–90.
34. Ayranci Ü, Yenilmez Ç. Eskişehir İlinde Birinci Basamak Sağlık Kurumlarında Verilen Ruh Sağlığı Hizmetlerinin Değerlendirilmesi. *Türk Psikiyatr Derg.* 2002;13(2):115–24.
35. Kitchener BA, Jorm AF. Mental health first aid training: a review of evaluation studies. *Aust N Z J Psychiatry.* 2006;40(1):6–8.
36. Link BG, Phelan JC. Labeling and stigma. In: Scheid TL, Brown TN editors. *In A Handbook for the Study of Mental Health: Social Contexts, Theories and Systems.* New York: Cambridge University Press; 2010. p. 571–87.